

Bibliography: Facility and Surgeon Volume

1. Jain S, Rosenbaum PR, Reiter JG, Ramadan OI, Hill AS, Silber JH, et al. Assessing the Ambulatory Surgery Center Volume-Outcome Association. *JAMA Surg.* 2024 Jan 24; e237161.
2. Barnes CL, Zhang X, Stronach BM, Haas DA. The Initial Impact of COVID-19 on Total Hip and Knee Arthroplasty. *J Arthroplasty.* 2021;36(7S):S56-S61.
3. Bouchard P, Demyttenaere S, Court O, Franco EL, Andalib A. Surgeon and hospital volume outcomes in bariatric surgery: a population-level study. *Surg Obes Relat Dis.* 2020;16(5):674-81.
4. Saulle R, Vecchi S, Cruciani F, Mitrova Z, Amato L, Davoli M. The combined effect of surgeon and hospital volume on health outcomes: a systematic review. *La Clinica Terapeutica.* 2019;170(2):e148-161.
5. Jolbäck P, Rolfson O, Cnudde P, Odin D, Malchau H, Lindahl H, et al. High annual surgeon volume reduces the risk of adverse events following primary total hip arthroplasty: a registry-based study of 12,100 cases in Western Sweden. *Acta Orthop.* 2019;90(2):153-158.
6. Mufarrih SH, Ghani MO, Martins RS, Qureshi NQ, Mufarrih SA, Malik AT, et al. Effect of hospital volume on outcomes of total hip arthroplasty: a systematic review and meta-analysis. *J Orthop Surg Res.* 2019;14(1):468.
7. Chou YY, Tung TC. Optimal Hospital and Surgeon Volume Thresholds to Improve 30-Day Readmission Rates, Costs, and Length of Stay for Total Hip Replacement. *J Arthroplasty.* 2019;34(9):1901-1908.
8. Wilson S, Marx RG, Pan TJ, Lyman S. Meaningful thresholds for the volume-outcome relationship in total knee arthroplasty. *J Bone Joint Surg Am.* 2016;98(20):1683-1690.
9. Zevin B, Aggarwal R, Grantcharov TP. Volume-outcome association in bariatric surgery: a systematic review. *Ann Surg.* 2012;256(1):60-71.
10. Lau RL, Perruccio AV, Gandhi R, Mahomed NN. The role of surgeon volume on patient outcome in total knee arthroplasty: a systematic review of the literature. *BMC Musculoskel Dis.* 2012;13(1):250.
11. Singh JA, Kwoh CK, Boudreau RM, Lee GC, Ibrahim SA. Hospital volume and surgical outcomes after elective hip/knee arthroplasty: a risk-adjusted analysis of a large regional database. *Arthritis Rheum.* 2011;63(8):2531-2539.
12. Birkmeyer NJO, Dimick JB, Share D, Hawasli A, English WJ, Genaw J, et al., Hospital complication rates with bariatric surgery in Michigan. *JAMA.* 2010;304(1):435-442.
13. Weller W, Hannan E. Relationship between provider volume and postoperative complications for bariatric procedures in New York State. *J Am Coll Surg.* 2006;202(5):753-761.
14. Ngyuen N, Paya M, Stevens C, Mavandadi S, Zainabadi K, Wilson S. The relationship between hospital volume and outcome in bariatric surgery at academic medical centers. *Ann Surg.* 2004;240(4):586-594.