

1. Jolbäck P, Rolfson O, Cnudde P, Odin D, Malchau H, Lindahl H, et al. High annual surgeon volume reduces the risk of adverse events following primary total hip arthroplasty: a registry-based study of 12,100 cases in Western Sweden. *Acta Orthop*. 2019;90(2):153-158.
2. Mufarrih SH, Ghani MO, Martins RS, Qureshi NQ, Mufarrih SA, Malik AT, et al. Effect of hospital volume on outcomes of total hip arthroplasty: a systematic review and meta-analysis. *J Orthop Surg Res*. 2019;14(1):468.
3. Wilson S, Marx RG, Pan TJ, Lyman S. Meaningful thresholds for the volume-outcome relationship in total knee arthroplasty. *J Bone Joint Surg Am*. 2016;98(20):1683-1690.
4. Lau RL, Perruccio AV, Gandhi R, Mahomed NN. The role of surgeon volume on patient outcome in total knee arthroplasty: a systematic review of the literature. *BMC Musculoskel Dis*. 2012;13(1):250.
5. Singh JA, Kwoh CK, Boudreau RM, Lee GC, Ibrahim SA. Hospital volume and surgical outcomes after elective hip/knee arthroplasty: a risk-adjusted analysis of a large regional database. *Arthritis Rheum*. 2011;63(8):2531-2539.